

NPS55-79-009

NAVAL POSTGRADUATE SCHOOL

Monterey, California



HEAVY TRAFFIC RESULTS
FOR SINGLE SERVER QUEUES WITH
DEPENDENT (EARMA) SERVICE
AND INTERARRIVAL TIMES

by

P. A. Jacobs

April 1979

Approved for public release; distribution unlimited.

NAVAL POSTGRADUATE SCHOOL
MONTEREY, CALIFORNIA

Rear Admiral T. F. Dedman
Superintendent

J. R. Borsting
Provost

Reproduction of all or part of this report is authorized.

This report was prepared by:

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NPS55-79-009	2. GOVT ACCESSION NO. AD-A070 086	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Heavy Traffic Results for Single Server Queues with Dependent (EARMA) Service and Interarrival Times		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) P. A. Jacobs		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, CA 93940		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Postgraduate School Monterey, CA 93940		10. PROGRAM ELEMENT PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE April 1979
		13. NUMBER OF PAGES 27
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Cross correlated interarrival and service times Multivariate exponential random variables Heavy traffic Weak convergence EARMA sequences		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Models are given for sequences of correlated exponential interarrival and service times for a single server queue. These multivariate exponential models are formed as probabilistic linear combinations of sequences of independent exponential random variables and are easy to generate on a computer. Limiting results for customer waiting time under heavy traffic conditions are obtained for these queues. Heavy traffic results are useful for analyzing the effect of correlated interarrival and service times in queues on		

20. ABSTRACT

such quantities as queue length and customer waiting time. They can also be used to check simulation results.

HEAVY TRAFFIC RESULTS FOR SINGLE SERVER QUEUES
WITH DEPENDENT (EARMA) SERVICE
AND INTERARRIVAL TIMES

by

P. A. Jacobs^{*}

Naval Postgraduate School
Monterey, California

0. ABSTRACT

Models are given for sequences of correlated exponential interarrival and service times for a single server queue. These multivariate exponential models are formed as probabilistic linear combinations of sequences of independent exponential random variables and are easy to generate on a computer. Limiting results for customer waiting time under heavy traffic conditions are obtained for these queues. Heavy traffic results are useful for analyzing the effect of correlated interarrival and service times in queues on such quantities as queue length and customer waiting time. They can also be used to check simulation results.

^{*}Support from NSF grants ENG-77-09020 and ENG-79-01438 and ONR contract NR-42-284 is gratefully acknowledged.

1. INTRODUCTION

Much of the work in queueing theory deals with models in which assumptions of independent service and interarrival times are made. One way to examine the effect of correlated interarrival and service times on measures of performance in queues is to consider models in which the interarrival and service times are correlated stationary sequences of random variables having given marginal distributions and compare results for these models with those for which the interarrival and service times are independent and identically distributed with the same marginal distributions.

In [3] Jacobs and Lewis introduced a scheme for generating sequences of dependent exponential random variables. The EARMA (1,1) (exponential mixed autoregressive moving average with both autoregression and moving average of order 1) model is defined as follows. Let $\{E_n\}$ be a sequence of independent exponential random variables with positive finite mean ν^{-1} . Let $\{J_n\}$ and $\{K_n\}$ be independent sequences of independent $\{0,1\}$ -random variables with $P\{J_n=1\} = 1 - \beta$ and $P\{K_n=1\} = 1 - \rho$ where $0 \leq \beta \leq 1$ and $0 \leq \rho < 1$ are given constants. We define

$$(1.1) \quad X_n = \beta E_n + J_n A_{n-1}, \quad n = 1, 2, \dots$$

where

$$(1.2) \quad A_n = \rho A_{n-1} + K_n E_n, \quad n = 1, 2, \dots$$

We assume that A_0 is an independent exponential random variable with mean ν^{-1} . Under this assumption $\{X_n\}$ is a stationary sequence of dependent exponential random variables with mean ν^{-1} .

The dependence is not Markovian in general and as was shown in [3]

$$(1.3) \quad \text{Corr}(X_1, X_{1+n}) = \rho^{n-1} (1-\beta) [\beta(1-\rho) + (1-\beta)\rho] .$$

Note that if $\beta = 1$ or $\beta = 0$ and $\rho = 0$, then $\{X_n\}$ is a sequence of independent exponential random variables.

The EARMA(1,1) process is formed as a probabilistic linear combination of independent random variables. This technique can be extended to form sequences of multivariate exponential random variables. This idea was briefly explored in Lewis and Shedler [7]. These models have the advantage that the marginal distribution and correlation structure of the sequence are specified separately. In section 2 we will give several ways in which to generate sequences of dependent multivariate exponential random variables from probabilistic linear combinations of independent exponentials. These stationary multivariate EARMA sequences of dependent exponentials can be used as interarrival and service times in queues. Results for measures of performance for these queues can then be compared to queues with stationary independent sequences of interarrival and service times having the same marginal (exponential) distributions. Since the correlation structure in the multivariate EARMA sequences is specified by parameters independent of the marginal distribution, any difference between results for queues with these multivariate interarrival and service sequences and the corresponding queues with independent interarrival and service sequences having the same marginal distributions will be due to the correlation structure.

Suppose that the interarrival and service times of a single server queue form stationary multivariate EARMA sequences

of dependent exponential random variables. Results of Loynes [9] give conditions for the existence of limiting distributions for the waiting time of a customer and the queue length. However, it seems to be difficult to obtain exact results even in the simplest queues with EARMA interarrival and service times (cf. Jacobs [4]).

As a result we turn our attention to the possibilities of obtaining limiting results of the heavy traffic type for queues with stationary multivariate EARMA sequences of dependent interarrival and service times. Results of this kind will indicate how dependence in queues introduced by the EARMA scheme affects quantities such as customer waiting time and queue length, at least under heavy traffic conditions. These limiting results can also be used to check simulations.

In this paper we will show that Kingman's [5] result for the equilibrium waiting time W in heavy traffic holds for queues having multivariate EARMA interarrival and service times; that is, we will show under conditions detailed in Section 3, that if the traffic intensity $\lambda\mu^{-1}$ of the queue is close to 1, then the distribution of $(1 - \lambda\mu^{-1})W$ is approximately exponential. The mean of this exponential distribution will in general be different from the case of independent interarrival and service times. Positive cross-correlation between the interarrival and service times will tend to decrease the mean; while positive correlation within the interarrival and service time sequences will tend to increase it.

In Section 3 we will state the heavy traffic result more precisely and present several examples of its application. Its formulation for multivariate EARMA queues is very dependent on the fact that the marginal distribution and correlation structure are specified independently of one another in EARMA formulations. In Section 4 we will give the proof of the heavy traffic result.

2. QUEUES WITH EARMA INTERARRIVAL AND SERVICE TIMES

We will consider a single server queue at which customers arrive and are served in the order of their arrival. The 0th customer arrives at $t = 0$ and finds the server free. The arrival time, service time, and waiting time (excluding service) of the n th customer are denoted respectively by $T_n = \sum_{k=1}^n X_k$, S_n , and W_n . In this section we will present some examples of the use of the EARMA scheme to model dependent interarrival and service times in queues.

(2.1) EXAMPLE. Let $\{X_n\}$ be an EARMA(1,1) process with positive finite mean λ^{-1} and parameters $0 \leq \beta_1 \leq 1$ and $0 \leq \rho_1 < 1$; let $\{S_n\}$ be an EARMA(1,1) process independent of $\{X_n\}$ with positive finite mean μ^{-1} and parameters $0 \leq \beta_2 \leq 1$ and $0 \leq \rho_2 < 1$. Note that if $\beta_1 = 1$, $\beta_2 = 1$, or $\beta_1 = 1$, $\beta_2 = 0$, $\rho_2 = 0$, or $\beta_1 = 0$, $\rho_1 = 0$, $\beta_2 = 1$, or $\beta_1 = 0$, $\rho_1 = 0$, $\beta_2 = 0$, $\rho_2 = 0$, then the queue reduces to an M/M/1 queue. Other special cases include queues with Poisson arrivals and EARMA(1,1) service times independent of the arrival process, and queues with EARMA(1,1) interarrival times and independent exponential service times.

The sequence $\{(X_n, S_n)\}$ is stationary and $\{X_n\}$ and $\{S_n\}$ are independent. However the X_n 's (respectively S_n 's) are positively correlated with

$$(2.2) \quad \text{Cov}(X_n, X_{n+k}) = \lambda^{-2} \rho_1^{k-1} (1-\beta_1) [\beta_1 (1-\rho_1) + (1-\beta_1) \rho_1]$$

(respectively

$$(2.3) \quad \text{Cov}(S_n, S_{n+k}) = \mu^{-2} \rho_2^{k-2} (1-\beta_2) [\beta_2 (1-\rho_2) + (1-\beta_2) \rho_2]$$

from (1.2).

Let \underline{F}_m^1 (respectively \underline{F}_m^2) be the σ -algebra generated by X_1, \dots, X_m (respectively S_1, \dots, S_m) and \underline{G}_{m+k}^1 (respectively \underline{G}_{m+k}^2) be the σ -algebra generated by $X_{m+k}, X_{m+k+1}, \dots$ (respectively $S_{m+k}, S_{m+k+1}, \dots$). Let $L^2(\underline{F}_m^i)$ (respectively $L^2(\underline{G}_{m+k}^i)$) denote the collection of real-valued functions having finite second moment that are measurable with respect to \underline{F}_m^i (respectively \underline{G}_{m+k}^i), $i = 1, 2$. If $f_i \in L^2(\underline{F}_m^i)$ and $g_i \in L^2(\underline{G}_{m+k}^i)$, $i = 1, 2$, then there is a k_0 independent of f_i, g_i , $i = 1, 2$, such that for $k \geq k_0$

$$\begin{aligned} (2.4) \quad & |E[(f_1 f_2)(g_1 g_2)] - E[f_1 f_2] E[g_1 g_2]| \\ &= |E[f_1 g_1] E[f_2 g_2] - E[f_1] E[f_2] E[g_1] E[g_2]| \\ &= |E[f_1 g_1] \{E[f_2 g_2] - E[f_2] E[g_2]\} \\ &\quad + E[f_2] E[g_2] \{E[f_1 g_1] - E[f_1] E[g_1]\}| \end{aligned}$$

$$\begin{aligned}
(2.5) \quad & \leq E[|f_1 g_1|] \leq \rho_2^{k/2} E[f_2^2]^{1/2} E[g_2^2]^{1/2} \\
& + E[|f_2|] E[|g_2|] \leq \rho_1^{k/2} E[f_1^2]^{1/2} E[g_1^2]^{1/2} \\
& \leq 5[\rho_1^{k/2} + \rho_2^{k/2}] E[(f_1 f_2)^2]^{1/2} E[(g_1 g_2)^2]^{1/2}
\end{aligned}$$

where (2.4) follows from the independent of $\{X_n\}$ and $\{S_n\}$ and (2.5) follows from (6.5) of Jacobs and Lewis [3]. Hence, if f (respectively g) has finite second moment and is measurable with respect to the product σ -algebra $\underline{F}_m^1 \times \underline{F}_m^2$ (respectively $\underline{G}_{m+k}^1 \times \underline{G}_{m+k}^2$), then for k sufficiently large

$$(2.6) \quad |E[fg] - E[f] E[g]| \leq 5E[f^2]^{1/2} E[g^2]^{1/2} [\rho_1^{k/2} + \rho_2^{k/2}].$$

Hence, the sequence $\{(X_n, S_n)\}$ is asymptotically uncorrelated in the sense of Rosenblatt [10] and is ϕ -mixing in the sense of Billingsley [1].

(2.7) EXAMPLE. In Lewis and Shedler [7] one of the schemes that is proposed for cross-correlating the arrival process and the service times is the following. Assume $\{X_n\}$ and $\{E_n\}$ are independent sequences of independent exponential random variables with positive finite means λ^{-1} and μ^{-1} respectively. The n th interarrival time is assumed to be X_n and hence the arrival process is Poisson with rate λ . Let the n th service time

$$(2.8) \quad S_n = \beta E_n + J_n(\lambda \mu^{-1} B_n)$$

where

$$(2.9) \quad B_n = \rho B_{n-1} + K_n X_n$$

and β , ρ , $\{J_n\}$ and $\{K_n\}$ are as in (1.1) and (1.2) so that $\{S_n\}$ and $\{X_n\}$ are now cross-correlated sequences. We will assume that B_0 has an exponential distribution with mean λ^{-1} .

Since E_n and B_n are independent, $\{S_n\}$ is a stationary sequence of exponential random variables with mean μ^{-1} . If $\beta = 1$, then the queue reduces to an M/M/1 queue. The S_n 's can be shown to be positively correlated with

$$(2.10) \quad \text{Cov}(S_n, S_{n+k}) = \mu^{-2} (1-\beta)^2 \rho^k, \quad k = 1, 2, \dots$$

Further, the n th service time S_n is positively correlated with the interarrival times X_n, X_{n-1}, \dots, X_1 with

$$(2.11) \quad \text{Cov}(S_n, X_{n-k}) = (\lambda\mu)^{-1} (1-\beta) (1-\rho) \rho^k$$

for $k = 0, \dots, n-1$.

Note that $(X_n, S_n, \dots, X_{n+k}, S_{n+k})$ is a function of $(B_{n-1}, X_n, E_n, J_n, K_n, X_{n+1}, E_{n+1}, J_{n+1}, K_{n+1}, \dots, X_{n+k}, E_{n+k}, J_{n+k}, K_{n+k})$. Hence, since the sequence $\{B_n\}$ is stationary so is the sequence $\{(X_n, S_n)\}$. Further, by arguments similar to those used in obtaining (6.5) of Jacobs and Lewis [3], if f and g are as in (2.6), then for k sufficiently large

$$(2.12) \quad |E[fg] - E[f] E[g]| \leq 5\rho^{k/2} E[f^2]^{1/2} E[g^2]^{1/2}$$

so that the sequence $\{(X_n, S_n)\}$ is ϕ -mixing.

There are many ways to cross-correlate the interarrival and service times using the EARMA scheme beyond those given above and in Lewis and Shedler [7]. We will present one more, somewhat more general than that of Example (2.7) in which the service times were coupled back over the previous interarrival times.

(2.13) EXAMPLE. Let $\{E_n\}$, $\{E_n\}$, and $\{C_n\}$ be independent sequences of independent exponential random variables having respective positive finite means λ^{-1} , μ^{-1} , and 1. Let

$$(2.14) \quad X_n = \beta E_n + J_n (\lambda^{-1} A_n)$$

$$(2.15) \quad S_n = \gamma E_n + I_n (\mu^{-1} A_n)$$

with

$$(2.16) \quad A_n = \rho A_{n-1} + K_n C_n$$

where β , ρ , $\{J_n\}$ and $\{K_n\}$ are as in (1.1) and (1.2) and $\{I_n\}$ is a sequence of independent binary random variables independent of everything with $P\{I_n=1\} = 1 - P\{I_n=0\} = (1-\gamma)$ for some fixed constant $0 \leq \gamma \leq 1$. Again we assume that A_0 is an independent random variable having an exponential distribution with mean 1. Then $\{(E_n, E_n, A_n, I_n, J_n)\}$ is a stationary sequence of random variables and, hence, the same is true of $\{(X_n, S_n)\}$.

If $\gamma = 1, \beta = 1$ or $\gamma = 1, \beta = 0, \rho = 0$ or $\gamma = 0, \beta = 1, \rho = 0$, then the queue reduces to an M/M/1 queue. If $\beta = 1, \rho = 0$ then the queue reduces to a special case of Example (2.7). Straightforward manipulation shows that in general the interarrival times X_n are positively correlated with

$$(2.17) \quad \text{Cov}(X_n, X_{n+k}) = \lambda^{-2} (1-\beta)^2 \rho^k, \quad k = 1, 2, \dots$$

The service times S_n are also positively correlated with

$$(2.18) \quad \text{Cov}(S_n, S_{n+k}) = \mu^{-2} (1-\gamma)^2 \rho^k, \quad k = 1, 2, \dots$$

The interarrival times and service times are positively cross-correlated with

$$(2.19) \quad \text{Cov}(X_n, S_{n+k}) = (1-\beta)(1-\gamma)(\lambda\mu)^{-1} \rho^k, \quad k = 0, 1, \dots;$$

and

$$(2.20) \quad \text{Cov}(X_{n+k}, S_n) = (1-\beta)(1-\gamma)(\lambda\mu)^{-1} \rho^k, \quad k = 0, 1, \dots$$

Finally, by the arguments used in obtaining (6.5) in Jacobs and Lewis [3], if f and g are as in (2.6), then for k sufficiently large

$$(2.21) \quad |E[fg] - E[f] E[g]| \leq 5\rho^{k/2} E[f^2]^{1/2} E[g^2]^{1/2}.$$

so that the sequence $\{(X_n, S_n)\}$ is ϕ -mixing.

3. A HEAVY TRAFFIC RESULT FOR EARMMA QUEUES

Let $\{X_n\}$ (respectively $\{S_n\}$) be the sequence of interarrival times (respectively service times) for a single server queue with first-in-first-out service discipline.

Put $U_n = S_n - X_n$ and assume that $\{U_n\}$ is a stationary sequence with $E[U_n] < 0$. Let W_n be the waiting time of the n th customer and $W = \sup_{n \geq 0} \sum_{k=1}^n U_k$ (with the convention that $\sum_{k=1}^0 = 0$). Loynes [9] has shown that if the sequence $\{U_n\}$ is metrically transitive, then the distribution of W_n tends in the limit to the distribution function of W . Since it is usually difficult to obtain the distribution of W analytically, particularly when the service and interarrival times are dependent, we will obtain a heavy traffic limiting result for it. In this section we will give the statement of the result as well as some examples of its use leaving the proof to the next section.

For each $\alpha > 0$ let Q_α be a single server queue with FIFO queueing discipline having stationary sequences of interarrival times $\{X_n(\alpha)\}$ with positive finite mean $\lambda(\alpha)^{-1}$ and service times $\{S_n(\alpha)\}$ with positive finite mean $\mu(\alpha)^{-1}$ such that, if $U_n(\alpha) = S_n(\alpha) - X_n(\alpha)$, then $\{U_n(\alpha)\}$ is a stationary sequence of random variables. Let

$$(3.1) \quad E[U_n(\alpha)] = v(\alpha) < 0; \text{Var}[U_n(\alpha)] = s^2(\alpha);$$

$$(3.2) \quad \sigma^2(\alpha) = \text{Var } U_1(\alpha) + 2 \sum_{n=1}^{\infty} \text{Cov}(U_1(\alpha), U_{1+n}(\alpha));$$

$$(3.3) \quad d(\alpha) = |v(\alpha)|/\sigma(\alpha) \quad \text{and} \quad n(\alpha) = d(\alpha)^{-2}.$$

We will assume the following:

(3.4) ASSUMPTION. There exists a distribution function G so that $\lim_{\alpha \rightarrow \infty} P\{[U_n(\alpha) - v(\alpha)]^2 \leq x\} = G(x)$ for all continuity points x of G ; $\lim_{\alpha \rightarrow \infty} v(\alpha) = 0$; and $0 < \lim_{\alpha \rightarrow \infty} \sigma^2(\alpha) = \sigma^2 < \infty$.

$$(3.5) \quad \text{ASSUMPTION.} \quad 0 < \lim_{\alpha \rightarrow \infty} \sigma^2(\alpha) = \sigma^2 < \infty.$$

(3.6) ASSUMPTION. If f (respectively g) is a function with finite second moment measurable with respect to the σ -algebra generated by $U_1(\alpha), \dots, U_m(\alpha)$ (respectively $U_{m+k}(\alpha), U_{m+k+1}(\alpha), \dots$), then

$$(3.7) \quad |E[fg] - E[f]E[g]| \leq \phi(k) E[f^2]^{1/2} E[g^2]^{1/2}$$

for some decreasing function ϕ bounded above by 1 not depending on α satisfying

$$(3.8) \quad \sum_{n=1}^{\infty} \phi(n) < \infty.$$

Assumption (3.4) concerns the convergence of the distribution of $U_n^2(\alpha) = [S_n(\alpha) - X_n(\alpha)]^2$ and its first two moments as $\alpha \rightarrow \infty$. Assumption (3.6) is roughly a condition that the sequences $\{U_n(\alpha), n=1,2,\dots\}$

be ϕ -mixing uniformly over all α . This assumption guarantees that the series in (3.2) converges and

$$\lim_{n \rightarrow \infty} \frac{1}{n} E \left[\left(\sum_{k=1}^n (U_k(\alpha) - v(\alpha)) \right)^2 \right] = \sigma^2(\alpha).$$

Assumption (3.5) concerns the convergence of this normalized variance term as $\alpha \rightarrow \infty$.

Put

$$Z_\alpha(t) = d(\alpha) \sigma(\alpha)^{-1} \sum_{k=1}^{[n(\alpha)t]} U_k(\alpha), \quad t \geq 0$$

where $[x]$ is the largest integer less than or equal to x .

Let $\mathbb{D}([0, \infty))$ be the collection of real-valued, right continuous functions on $[0, \infty)$ which have left limits everywhere endowed with the metric given in Lindvall [8].

(3.9) THEOREM. If (3.4)-(3.6) are satisfied, then the process $Z_\alpha = \{Z_\alpha(t); t \geq 0\}$ converges weakly in $\mathbb{D}([0, \infty))$ to a Brownian motion with negative unit drift.

The proof of this result will be given in the next section.

Let

$$W(\alpha) = \sup_{n \geq 0} \sum_{k=1}^n U_k(\alpha).$$

Then it follows from Theorem (3.9) that

$$\lim_{\alpha \rightarrow \infty} P \left\{ \frac{d(\alpha)}{\sigma(\alpha)} W(\alpha) \leq x \right\} = \lim_{\alpha \rightarrow \infty} P \left\{ \sup_{t \geq 0} Z_{\alpha}(t) \leq x \right\}$$

$$= 1 - \exp\{-2x\}$$

where the last equality follows from the continuous mapping theorem (cf. Billingsley [1]). Thus we have the following result.

(3.10) THEOREM. If (3.4)-(3.6) hold, then

$$\lim_{x \rightarrow \infty} P\{|v(x)| W(\alpha) \leq x\} = 1 - \exp\{-2\sigma^{-2}x\}.$$

Hence, if the traffic intensity $\lambda\mu^{-1}$ of a queue is less than but close to 1 and $\{U_n\}$ is a stationary sequence satisfying (3.7) and (3.8), then the distribution of W is approximately exponential with mean $-\frac{1}{2}\sigma^2 E[U_1]^{-1}$ where

$$\sigma^2 = \text{Var}[U_1] + 2 \sum_{n=1}^{\infty} \text{Cov}(U_1, U_{1+n}).$$

Thus positive cross correlation between the interarrival and service times will tend to decrease the average mean waiting time; while positive correlation within the interarrival time or service time sequence will tend to increase the mean waiting time.

(2.11) EXAMPLE. For each $\alpha > 0$, let Q_α be a queue as in Example (2.1) with $\{X_n(\alpha)\}$ (respectively $\{S_n(\alpha)\}$) forming an EARMA(1,1) process with parameters $\lambda(\alpha)$, β_1 and ρ_1 (respectively $\mu(\alpha)$, β_2 and ρ_2). Assume $\lambda(\alpha) < \mu(\alpha)$ for each α , but $\lim_{\alpha \rightarrow \infty} \lambda(\alpha) = \lim_{\alpha \rightarrow \infty} \mu(\alpha) = m$; then hypotheses (3.4)-(3.6) are satisfied and Theorem (3.10) holds with

$$\sigma^2 = 2m^{-2} \{1 + (1-\rho_1)^{-1} (1-\beta_1) [(1-\rho_1)\beta_1 + (1-\beta_1)\rho_1] + (1-\rho_2)^{-1} (1-\beta_2) [(1-\rho_2)\beta_2 + (1-\beta_2)\rho_2]\}.$$

Hence, if the traffic intensity $\lambda\mu^{-1}$ of a queue with independent EARMA(1,1) services and interarrival times is less than but close to 1, then the distribution of the equilibrium waiting time W is approximately exponential with mean $-\frac{1}{2} \sigma^2 E[U_1]$ where $E[U_1] = \mu^{-1} - \lambda^{-1}$ and σ^2 is given by

$$\sigma^2 = \lambda^{-2} \{1 + 2(1-\rho_1)^{-1} (1-\beta_1) [(1-\rho_1)\beta_1 + (1-\beta_1)\rho_2]\} + \mu^{-2} \{1 + 2(1-\rho_2)^{-1} (1-\beta_2) [(1-\rho_2)\beta_2 + (1-\beta_2)\rho_2]\}.$$

Thus the dependence in this queue tends to increase the mean customer waiting time at least in close to heavy traffic conditions.

(3.12) EXAMPLE. For each $\alpha > 0$, let Q_α be a queue as in Example (2.7) with parameters $E[X_n(\alpha)] = \lambda(\alpha)^{-1}$, β , and ρ and $E[S_n(\alpha)] = \mu(\alpha)^{-1}$. Again we will assume $\lambda(\alpha) < \mu(\alpha)$ for each α but $\lim_{\alpha \rightarrow \infty} \lambda(\alpha) = \lambda = \lim_{\alpha \rightarrow \infty} \mu(\alpha) = \mu = m$. Hence (3.4)-(3.6) are satisfied and Theorem (3.10) holds with

$$\begin{aligned}\sigma^2 &= \lambda^{-2} + \mu^{-2} - 2(1-\beta)(1-\rho)(\lambda\mu)^{-1} \\ &\quad + 2\rho(1-\rho)^{-1} [(1-\beta)^2\mu^{-2} - (1-\beta)(1-\rho)(\lambda\mu)^{-1}] \\ &= \lambda^{-2} + \mu^{-2} [1 + 2(1-\beta)^2\rho(1-\rho)^{-1}] - 2(\lambda\mu)^{-1}(1-\beta) \\ &= 2m^{-2} [1 + (1-\beta)^2\rho(1-\rho)^{-1} - (1-\beta)] .\end{aligned}$$

The quantity $\sigma^2 - \sigma_{\text{ind}}^2$ can be either positive or negative where $\sigma_{\text{ind}}^2 = \lambda^{-2} + \mu^{-2} = 2m^{-2}$. It is negative if $\rho < (2-\beta)^{-1}$ and nonnegative otherwise. Hence, if ρ is sufficiently less than $(2-\beta)^{-1}$, then the average customer waiting time in the queue will tend to be less than the waiting time in the independent case at least in close to heavy traffic conditions. It will tend to be greater if ρ is sufficiently greater than $(2-\beta)^{-1}$.

(3.13) EXAMPLE. For each $\alpha > 0$, let Q_α be a queue as in Example (2.13) with parameters $\lambda(\alpha)$, $\mu(\alpha)$, ρ , β , and γ . Assume $\lim_{\alpha \rightarrow \infty} \lambda(\alpha) = \lambda = \lim_{\alpha \rightarrow \infty} \mu(\alpha) = \mu = m$ as before. Then (3.4)-(3.6) are satisfied and Theorem (3.10) holds with

$$\begin{aligned}
\sigma^2 &= \mu^{-2} + \lambda^{-2} - 2(\lambda\mu)^{-1} (1-\beta)(1-\gamma) \\
&\quad + 2\rho(1-\rho)^{-1} \{\mu^{-1}(1-\gamma) - \lambda^{-1}(1-\beta)\}^2 \\
&= 2m^{-2} [1 - (1-\beta)(1-\gamma) + \rho(1-\rho)^{-1} \{(\beta-\gamma)\}^2] .
\end{aligned}$$

The difference $\sigma^2 - \sigma_{\text{ind}}^2$ can be either positive or negative; ($\sigma_{\text{ind}}^2 = \lambda^{-2} + \mu^{-2} = 2m^{-2}$). The difference will be negative if $\rho < (1-\beta)(1-\gamma)/[(1-\beta)(1-\gamma) + (\beta-\gamma)^2]$ and nonnegative otherwise. Hence under close to heavy traffic conditions, the average customer waiting time will tend to be less than the independent case if $(1-\beta)(1-\gamma)/[(1-\beta)(1-\gamma) + (\beta-\gamma)^2]$ is sufficiently greater than ρ . It will be greater than the independent case if the same quantity is sufficiently less than ρ .

4. PROOF OF THE MAIN RESULT

In this section we will give a proof of Theorem (3.9). We will assume that (3.4)-(3.6) are satisfied. Let

$$Y_{\alpha}(t) = \frac{d(\alpha)}{\sigma(\alpha)} \sum_{k=1}^{[n(\alpha)t]} [U_k(\alpha) - v(\alpha)] .$$

By Theorem 3' of Lindvall [8] and the continuous mapping theorem, Theorem (3.9) will be proved if for each $a > 0$ we can show the following result.

(4.1) THEOREM. The process $\{Y_{\alpha}(t); 0 \leq t \leq a\}$ converges weakly on $\mathbb{D}([0,a])$ to Brownian motion.

($\mathbb{D}([0,a])$ is the space of real-valued right continuous functions on $[0,a]$ with left-hand limits endowed with the Skorohod J_1 topology.)

We will prove Theorem (4.1) for $a = 1$ by a series of lemmas. The proof for arbitrary $a > 0$ is similar.

(4.2) LEMMA. The process $\{Y_{\alpha}(t); 0 \leq t \leq 1\}$ has asymptotically independent increments.

Proof. Let $0 \leq s_1 \leq t_1 < s_2 \leq t_2 < \dots < s_r \leq t_r \leq 1$. Let C_i be the event $\{Y_{\alpha}(t_i) - Y_{\alpha}(s_i) \leq x_i\}$ for real x_i . Then C_i is an event in the σ -algebra generated by

$U_{[n(\alpha)s_i]+1}, \dots, U_{[n(\alpha)t_i]}$. If δ is the smallest difference $s_i - t_{i-1}$, then $[n(\alpha)s_i] + 1 - [n(\alpha)t_{i-1}] \geq [n(\alpha)\delta]$. By (3.7)

$$|P(\prod_{i=1}^r C_i) - \prod_{i=1}^r P(C_i)| \leq r\phi([n(\alpha)\delta]) .$$

Since $\delta > 0$, $r\phi([n(\alpha)\delta])$ tends to zero as $\alpha \rightarrow \infty$ and the result follows.

In the following results, let $D_i(\alpha) = U_i(\alpha) - v(\alpha)$ and $V_m(\alpha) = \sum_{i=1}^m D_i(\alpha)$. By (3.4)-(3.6) the following result holds.

(4.3) LEMMA.

$$a) \quad \frac{1}{k} E[V_k(\alpha)^2] \leq 2 \sum_{j=0}^{\infty} |E[D_1(\alpha) D_{1+j}(\alpha)]|;$$

$$b) \quad \lim_{\alpha \rightarrow \infty} n(\alpha)^{-1} E[V_{n(\alpha)}(\alpha)^2] = \sigma^2 ;$$

$$c) \quad \text{If } D_1 \text{ is bounded by } C, \text{ then } E[V_{n(\alpha)}(\alpha)^4] \leq K_{\phi} C^4 n(\alpha)^2$$

where K_{ϕ} depends on ϕ alone.

(4.4) LEMMA. The sequence of random variables $\{n(\alpha)^{-1}(V_{n(\alpha)}(\alpha))^2; \alpha \geq 0\}$ is uniformly integrable.

Proof. By (3.4), $\{D_1^2(\alpha); \alpha \geq 0\}$ is uniformly integrable. The remainder of the proof is very similar to that of (20.48) on page 176 of Billingsley [1] and will be omitted.

(4.5) LEMMA. For each positive ε there exists a γ , $\gamma > 0$ and an integer n_0 such that $n(\alpha) \geq n_0$ implies

$$P\left\{\max_{i \leq n(\alpha)} |V_i(\alpha)| \geq \gamma \sqrt{n(\alpha)}\right\} \leq \frac{\varepsilon}{\gamma^2}.$$

Proof. Put $C_i^*(\alpha) = \sum_{j=1}^i |D_j(\alpha)|$. By (3.4), there exists an increasing sequence of integers m_i such that $n(\alpha) \geq m_i$ implies $P\{|D_1(\alpha)| \geq \gamma \sqrt{n(\alpha)}/i^2\} \leq (\gamma n(\alpha) i^2)^{-1}$ for each positive γ . Let $p(\alpha) = i$ if $m_i \leq n(\alpha) < m_{i+1}$; then $p(\alpha)$ tends to infinity as $\alpha \rightarrow \infty$ but

$$(4.6) \quad \lim_{\alpha \rightarrow \infty} n(\alpha) P\{C_{p(\alpha)}^*(\alpha) \geq \gamma \sqrt{n(\alpha)}\}$$

$$\leq \lim_{\alpha \rightarrow \infty} n(\alpha) p(\alpha) P\{|D_1(\alpha)| \geq \gamma \sqrt{n(\alpha)} p(\alpha)^{-1}\}$$

$$\leq \lim_{\alpha \rightarrow \infty} n(\alpha) p(\alpha) P\{|D_1(\alpha)| \geq \gamma \sqrt{n(\alpha)} p(\alpha)^{-2}\}$$

$$\leq \lim_{\alpha \rightarrow \infty} n(\alpha) p(\alpha) [\gamma n(\alpha) p(\alpha)^2]^{-1} = 0.$$

We can further choose $p(\alpha)$ so that $p(\alpha) \leq n(\alpha)$. Fix $\varepsilon > 0$. By the previous lemma we can choose γ so that $\gamma > \sigma$ and

$$(4.7) \quad P\{|V_{k(\alpha)}(\alpha)| > \gamma \sqrt{k(\alpha)}\} < \varepsilon \gamma^{-2}$$

for all α . Let

$$\Lambda_i(\alpha) = \{\max_{j < i} |V_j(\alpha)| < 3\gamma \sqrt{n(\alpha)} \leq |V_i(\alpha)|\};$$

then

$$\begin{aligned} & P\{\max_{i \leq n(\alpha)} |V_i(\alpha)| \geq 3\gamma \sqrt{n(\alpha)}\} \\ & \leq P\{|V_{n(\alpha)}(\alpha)| \geq \gamma \sqrt{n(\alpha)}\} \\ & \quad + \sum_{i=1}^{n(\alpha)-1} P(\Lambda_i(\alpha) \cap \{|V_{n(\alpha)} - V_i(\alpha)| \geq 2\gamma \sqrt{n(\alpha)}\}) . \end{aligned}$$

The summation is bounded above by

$$\begin{aligned} & \sum_{i=1}^{n(\alpha)-p(\alpha)-1} P\{|V_i(\alpha) - V_{i+p(\alpha)}(\alpha)| \geq \gamma \sqrt{n(\alpha)}\} \\ & + \sum_{i=1}^{n(\alpha)-p(\alpha)-1} P(\Lambda_i(\alpha) \cap \{|V_{n(\alpha)}(\alpha) - V_{i+p(\alpha)}(\alpha)| > \gamma \sqrt{n(\alpha)}\}) \\ & + \sum_{i=n(\alpha)-p(\alpha)}^{n(\alpha)-1} P\{|V_{n(\alpha)}(\alpha) - V_i(\alpha)| \geq \gamma \sqrt{n(\alpha)}\} . \end{aligned}$$

Each term in the first and third of these sums is bounded above by $P\{C_{p(\alpha)}^*(\alpha) \geq \gamma \sqrt{n(\alpha)}\}$. Since $\Lambda_i(\alpha)$ is measurable with respect to the σ -algebra generated by $U_1(\alpha), \dots, U_i(\alpha)$ by (3.7)

$$P\{ \max_{i \leq n(\alpha)} |V_i(\alpha)| \geq 3\gamma \sqrt{n(\alpha)} \}$$

$$\leq P\{|V_{n(\alpha)}(\alpha)| \geq \gamma \sqrt{n(\alpha)}\}$$

$$+ [n(\alpha) + p(\alpha)] P\{C_{p(\alpha)}^*(\alpha) \geq \gamma \sqrt{n(\alpha)}\}$$

$$+ \sum_{i=1}^{n(\alpha)-p(\alpha)-1} P(\Lambda_i(\alpha)) [P\{|V_{n(\alpha)}(\alpha) - V_{i+p(\alpha)}(\alpha)| \geq \gamma \sqrt{n(\alpha)}\} + \phi(p(\alpha))]$$

$$\leq 2\varepsilon\gamma^{-2} + \phi(p(\alpha)) + [n(\alpha) + p(\alpha)] P\{C_{p(\alpha)}^*(\alpha) \geq \gamma \sqrt{n(\alpha)}\}$$

by (4.7) and the fact that the $\Lambda_i(\alpha)$ are disjoint. Hence by (4.6) and (3.8) for α large

$$P\{ \max_{i \leq n(\alpha)} |V_i(\alpha)| \geq 3\gamma \sqrt{n(\alpha)} \} \leq 3\varepsilon\gamma^{-2}.$$

PROOF OF THEOREM (4.1). For $a = 1$: By Lemma (4.5) the sequence $\{Y_\alpha\}$ is tight on $\mathbb{D}[0,1]$. Further $\lim_{\alpha \rightarrow \infty} E[Y_\alpha(t)] = 0$ and

$$\lim_{\alpha \rightarrow \infty} E[Y_\alpha(t)^2] = \lim_{\alpha \rightarrow \infty} (\sqrt{n(\alpha)} \sigma(\alpha))^{-2} E \left[\left(\sum_{k=1}^{[n(\alpha)t]} D_k(\alpha) \right)^2 \right] = t$$

by (3.5). Since $Y_\alpha(t)^2 = [\sigma^2(\alpha) n(\alpha)]^{-1} V_{[n(\alpha)t]}(\alpha)^2$,

$\{Y_\alpha^2(t)\}$ is uniformly integrable for each t by Lemma (4.4).

By Lemma (4.2) $\{Y_\alpha(t); 0 \leq t \leq 1\}$ has asymptotically independent increments. The result now follows from Theorem (19.2) of Billingsley [1].

5. EXTENSIONS

This paper presented a heavy traffic result for the limiting distribution of the customer waiting time in single server queues with dependent interarrival and service times. There are many different ways in which to model dependent interarrival and service times in queues using the EARMA scheme. Some other ways are given in Lewis and Shedler [7]. Properties of these multivariate exponential models will be given elsewhere.

Work is currently going on in the simulation of these queues. Some simulation results have been reported for the queue of Example (2.7) with $\beta = 0.5$, $\mu = 4$ and various values of ρ and λ in Boonsong [2]. They seem to indicate that the heavy traffic result approximates the average customer waiting time fairly well for a traffic intensity of .9 and quite well for a traffic intensity of .95. However, the distribution of W in a simulation with $\rho = 0.25$ and traffic intensity .995 shows a large amount of underdispersion relative to an exponential distribution. Thus, it seems that the convergence of the waiting time distribution is very slow.

It is expected that further simulations will give more indication of the validity of using the heavy traffic result to approximate the average customer waiting time, or quantiles of the waiting time distribution under less than heavy traffic conditions. These results will be reported on elsewhere.

6. REFERENCES

- [1] BILLINGSLEY, P. (1968). Convergence of Probability Measures. Wiley, New York.
- [2] BOONSONG, P. (1978). A Simulation Study of a Class of First-Come First-Served Queues with EARMA Correlation Structure, Master Thesis, Naval Postgraduate School, Monterey, California, September.
- [3] JACOBS, P. A. and LEWIS, P.A.W. (1977). A mixed auto-regressive-moving average exponential sequence and point process (EARMA 1,1). Adv. Appl. Prob. 9, 87-104.
- [4] JACOBS, P. A. (1978). A cyclic queueing network with dependent exponential service times. J. Appl. Prob. 15, 573-589.
- [5] KINGMAN, J. F. C. (1962). On queues in heavy traffic. J. Roy. Statist. B 24, 383-392.
- [6] LEWIS, P. A. W. and SHEDLER, G. S. (1977). Analysis and modelling of point processes in computer systems. Bull. Int. Statist. Inst. 47, 193-210.
- [7] LEWIS, P.A.W. and SHEDLER, G.S. (1977). Analysis and modelling of point processes in computer systems. Naval Postgraduate School Technical Report NPS55-77-38.
- [8] LINDVALL, T. (1973). Weak convergence of probability measures and random functions in the function space $D[0, \infty)$. J. Appl. Prob. 10, 109-121.
- [9] LOYNES, R.M. (1962). The stability of a queue with non-independent interarrival and services. Proc. Camb. Philos. Soc. 58, 497-520.
- [10] ROSENBLATT, M. (1971). Markov Processes. Structure and Asymptotic Behavior. Springer-Verlag, Berlin.

DISTRIBUTION LIST

	No. of Copies
Defense Documentation Center Cameron Station Alexandria, VA 22314	2
Technical Information Division Naval Research Laboratory Washington, D.C. 20375	1
Library Code 0212 Naval Postgraduate School Monterey, CA 93940	2
Library Code 55 Naval Postgraduate School Monterey, CA 93940	1
Dean of Research Code 012 Naval Postgraduate School Monterey, CA 93940	1
Statistics and Probability Program Office of Naval Research Code 436 Arlington, VA 22217	1
Library Naval Ocean Systems Ctr San Diego, CA 92152	1
Naval Electronic Systems Command Navelex 320 National Center No. 1 Arlington, VA 20360	1
Director Naval Research Lab Attn: Library (ONRL) Code 2029 Washington, D.C. 20375	1
Office of Naval Research San Francisco Area Office 760 Market Street San Francisco, CA 94102	1

	No. of Copies
Prof. T. W. Anderson Department of Statistics Stanford University Stanford, CA 94305	1
Professor F. J. Anscombe Department of Statistics Yale University New Haven, CT 06520	1
Professor C. R. Baker Department of Statistics University of North Carolina Chapel Hill, N.C. 37512	1
Prof. R. E. Barlow Operations Research Ctr College of Engineering University of California Berkeley, CA 94720	1
Prof. R. E. Bechhofer Department of Operations Research Cornell University Ithaca, New York 14850	1
Mr. C. N. Bennett Naval Coastal Systems Lab Code P761 Panama City, FL 32401	1
Prof. N. J. Bershad School of Engineering University of California Irvine, CA 92664	1
Prof. L. N. Bhat Comput Sci/Oper Res Ctr Southern Methodist Univ. Dallas, TX 75275	1
P. J. Bickel Department of Statistics University of California Berkeley, CA 94720	1
Prof. W. R. Blischke Dept. of Quantitative Business Analysis Univ. of Southern California Los Angeles, CA 90007	1

	No. of Copies
Prof. H. W. Block Dept. of Mathematics University of Pittsburgh Pittsburgh, PA 15260	1
Prof. Peter Bloomfield Dept. of Statistics Princeton University Princeton, N.J. 08540	1
Prof. Joseph Blum Dept. of Math., Stat. and Com. Sci. The American University Washington, D.C. 20016	1
J. E. Boyer, Jr. Dept. of Statistics Southern Methodist University Dallas, TX 75275	1
Dr. William Brogan National Science Foundation Systems Theory and Applications Program Eng. Div. 18th and "G", N.W. Washington, D.C. 20550	1
Prof. Ted Brown Dept. of Computer Science Queens College Flushing, N.Y. 11367	1
Dr. J. Chandra U. S. Army Research P.O. Box 12211 Research Triangle Park North Carolina 27706	1
Prof. H. Chernoff Dept. of Mathematics Mass. Inst. Tech. Cambridge, MA 02139	1
Prof. E. Çinlar Dept. of Industrial Engineering Northwestern University Evanston, ILL 60201	1
Dr. D. R. Cox Dept. of Mathematics Imperial College London SW7 2BZ, ENGLAND	1

No. of Copies

Prof. C. Derman Dept. of Civil Eng. and Eng. Mech. Columbia University New York, New York 10027	1
Prof. R. L. Disney Virginia Poly. Inst. and State Univ. Dept. of Ind. Eng. Blacksburg, VA 24061	1
Mr. J. Dowling Defense Logistics Studies Information Exchange Army Logistics Management Center Fort Lee, VA 20390	1
Dr. M. J. Fischer Defense Communications Agency 1860 Wiehle Ave. Reston, VA 22070	1
Prof. George S. Fishman Curr. in OR and Sys Anal. Univ. of North Carolina Phillips Annex Chapel Hill, N.C. 20742	1
Prof. Larry George Ind. Eng. Dept. Texas A&M University College Station, TX 77843	1
Mr. Gene H. Gleissner Appl. Math. Lab. David Taylor Naval Ship Research and Devel. Ctr Bethesda, MD 20084	1
Dr. R. Gnanadesikan Bell Telephone Lab. Murray Hill, N.J.	1
Dr. A. J. Goldman, Chief OR Div. 205.02, Admin. A428 U.S. Dept. of Commerce Washington, D.C. 20234	1
Prof. S. S. Gupta Dept. of Statistics Purdue University Lafayette, IN 47907	1

	No. of Copies
Prof. D. L. Hanson Dept. of Math. Sci. State University of New York, Binghamton, NY 13901	1
Prof. F. J. Harris Dept. of Elec. Eng. San Diego State Univ. San Diego, CA 92182	1
Prof. L. H. Herbach Dept. of Oper. Res. and Sys. Anal. Polytechnic Inst. of N.Y. Brooklyn, NY 11201	1
Prof. M. J. Hinich Dept. of Economics Virginia Polytechnic Inst. and State Univ. Blacksburg, VA 24061	1
Prof. W. M. Hirsch Inst. of Math. Sci. New York Univ. New York, N.Y. 19453	1
Prof. D. L. Iglehart Dept. of Oper. Res. Stanford Univ. Stanford, CA 94350	1
Prof. J. B. Kadane Dept. of Statistics Carnegie-Mellon Pittsburgh, PA 15213	1
Dr. H. Kobayashi IBM Yorktown Heights, N.Y. 10598	1
Dr. Guy Latouche Laboratoire d'Informatique Theorique Universite Libre de Bruxelles Boulevard du Triomphe 1050 Bruxelles, BELGIUM	1
Dr. Richard Lau, Director Office of Naval Research Branch Office 1030 Green Street Pasadena, CA 91101	1

	No. of Copies
Prof. N. Leadbetter Dept. of Statistics University of North Carolina Chapel Hill, N.C. 27415	1
Dr. John Lehoczky Dept. of Statistics Carnegie-Mellon Univ. Pittsburgh, PA 15213	1
Dr. A. Lemoine 1020 Guinda St. Palo Alto, CA 94301	1
Prof. R. S. Levenworth Dept. of Ind. and Sys. Eng. Univ. of Florida Gainesville, FL 32611	1
Prof. G. Lieberman Dept. of Oper. Res. Stanford University Stanford, CA 94305	1
Dr. J. MacQueen University of California Los Angeles, CA 90024	1
Dr. N. R. Mann Biomathematics Dept. School of Public Health Univ. of California Los Angeles, CA 90024	1
Dr. W. H. Marlow Program in Logistics George Washington University 707 22nd Street, N.W. Washington, D.C. 20037	1
Dr. M. Mazumdar Math. Dept. Westinghouse Labs Churchill Boro Pittsburgh, PA 15235	1
Dr. Masao Mori Dept. of Inf. Eng. Ibaraki Univ. Nakanarusawa-Machi 4-12-1 Hitachi, JAPAN	1

	No. of Copies
Prof. J. A. Muckstadt Dept. of Oper. Res. Cornell University Ithaca, N.Y. 19850	1
Dr. Janet M. Myhre Inst. Dec. Sci. for Bus. and Pub. Pol. Claremont Men's College Claremont, CA 91711	1
Prof. Marcel Neuts University of Delaware Dept. of Mathematics Newark, Delaware 19711	1
W. R. Nunn Center for Naval Analyses 1401 Wilson Blvd Arlington, VA 22209	1
B. S. Orleans Naval Sea Systems Command (Sea 03F), Rm 10SC8 Arlington, VA 20360	1
Prof. D. B. Owen Dept. of Statistics Southern Methodist Univ. Dallas, TX 75222	1
Prof. B. Pagurek Dept. of Sys Eng and Com Sci Carleton University Ottawa, Ont K1S 5B6 Canada	1
Prof. E. Parzen Statistics Dept. Texas A&M College Station, TX 77843	1
Prof. S. L. Phoenix Sibley School of Mech. and Aero. Eng. Cornell University Ithaca, N.Y. 14850	1
Prof. M. L. Puri Dept. of Mathematics P.O. Box F Indiana Univ. Foundation Bloomington, IN 47401	1

Prof. M. Rosenblatt Dept. of Mathematics Univ. of California San Diego, CA 92093	1
Prof. S. M. Ross College of Engineering Univ. of California Berkeley, CA 94720	1
M. N. Rothkopf Xerox Palo Alto Research Ctr 333 Coyote Hill Road Palo Alto, CA 94304	1
Robert Sargent Dept. of Ind. Eng. and Oper. Res. Syracuse Univ. Syracuse, N.Y. 13210	1
Prof. L. L. Scharf Jr. Dept. of Elec. Eng. Colorado State University Ft. Collins, CO 89521	1
Dr. Paul Schwitzer Graduate School of Management Univ. of Rochester Rochester, N.Y. 14627	1
Prof. D. G. Shimshak Univ. of Massachusetts Harbor Campus Boston, MA 02125	1
Prof. D. C. Siegmund Dept. of Statistics Stanford Univ. Stanford, CA 94305	1
Prof. W. L. Smith Dept. of Statistics Univ. of North Carolina Chapel Hill, N.C. 27514	1
Prof. J. G. Shanthikumar Dept. of Ind. Eng. University of Toronto Toronto, Ont. M5S 1R4 CANADA	1

	No. of Copies
Prof. J. R. Thompson Dept. of Math. Sci. Rice University Houston, TX 77001	1
Prof. J. L. Tukey Dept. of Statistics Princeton Univ. Princeton, N.J. 98540	1
Daniel H. Wagner Station Square One Paoli, PA 19301	1
Prof. Grace Wahba Dept. of Statistics Univ. of Wisconsin Madison, WI 53706	1
Prof. G. S. Watson Dept. of Statistics Princeton Univ. Princeton, N.J. 08540	1
Prof. Roy Welsch M.I.T. Sloan School of Management Cambridge, MA 02139	1
Naval Postgraduate School Monterey, CA. 93940	
Attn: Prof. D.P. Gaver, Code 55Gv	1
Prof. P.A. Jacobs, Code 55Jc	20
R. J. Stampfel, Code 55	1
Prof. R. A. Bradley Dept. of Statistics Florida State University Tallahassee, FL 32306	1

U-187,377

Naval Postgraduate School.
NPS-55-79-009.

30 MAY 97
6 NOV 90

33430
36471

U187377

DUDLEY KNOX LIBRARY - RESEARCH REPORTS



5 6853 01062193 1

0187377